

Effect Of Thiamine Pyrophosphate as an Adjunct to Metformin in Type 2 Diabetes: A Randomized, Controlled Pilot Clinical Trial

Melchor Alpízar Salazar^{1*}, María Fernanda Alpízar Sanchez¹, Magalie Manuell Retana², Eliane Durán Manuell², Carlos Jiménez Collado², Mónica Leticia Vallejo González¹, Miguel Alejandro Trejo¹, Priscila Judith Torres Granados¹, Jesús Manuel De Aldecoa Castillo¹ and Rohini Bala¹

¹Centro Especializado en Diabetes, Obesidad y Prevención de Enfermedades Cardiovasculares (CEDOPEC), Mexico

²Laboratorios Manuell, S A, Mexico City, Mexico.

***Corresponding author:** Melchor Alpízar Salazar, PhD, FACE, Centro Especializado en Diabetes, Obesidad y Prevención de Enfermedades Cardiovasculares (CEDOPEC), Mexico

ARTICLE INFO

Received: 📅 October 01, 2026

Published: 📅 October 22, 2026

Citation: Melchor Alpízar Salazar, María Fernanda Alpízar Sanchez, Magalie Manuell Retana, Eliane Durán Manuell, Carlos Jiménez Collado, Mónica Leticia Vallejo González, Miguel Alejandro Trejo, Priscila Judith Torres Granados, Jesús Manuel De Aldecoa Castillo and Rohini Bala. Effect Of Thiamine Pyrophosphate as an Adjunct to Metformin in Type 2 Diabetes: A Randomized, Controlled Pilot Clinical Trial. Biomed J Sci & Tech Res 67(2)-2026. BJSTR. MS.ID.010455.

ABSTRACT

Background: Type 2 diabetes mellitus (T2DM) is a chronic metabolic disorder characterized by persistent hyperglycemia and an increased risk of cardiovascular and microvascular complications. Functional thiamine deficiency and altered activity of thiamine-dependent enzymes have been implicated in the metabolic dysregulation associated with diabetes. Thiamine pyrophosphate (TPP), the biologically active form of vitamin B1, plays a central role in mitochondrial glucose metabolism and may contribute to improved metabolic control.

Methods: A randomized, placebo-controlled pilot clinical trial was conducted to evaluate the effect of TPP as an adjunct treatment to metformin in patients with T2DM with inadequate glycaemic control. Twenty-seven participants receiving stable metformin therapy were randomly assigned to receive weekly intravenous TPP or placebo for 12 weeks. Anthropometric, hemodynamic, and biochemical variables were assessed, including glycated haemoglobin (HbA1c), fasting glucose, and lipid profile. Inflammatory markers were also analysed. Both within-group and between-group statistical analyses were performed.

Results: Twenty-seven patients were included in the final analysis. The intervention group showed a significant reduction in HbA1c (-7.4%, $p = 0.003$) and systolic blood pressure (-5.3%, $p = 0.032$). Other metabolic and inflammatory parameters showed favourable trends without reaching statistical significance. The intervention was well tolerated, and no serious adverse events were reported.

Conclusion: Adjunctive therapy with thiamine pyrophosphate demonstrated promising effects on glycaemic control and hemodynamic parameters in patients with T2DM treated with metformin. These findings support the need for larger randomized clinical trials to confirm the potential therapeutic role of TPP in diabetes management.

Trial Registration: ClinicalTrials.gov (NCT04053621).

Keywords: Type 2 diabetes mellitus; Thiamine pyrophosphate; Metformin; Adjunctive Therapy; Glycaemic Control; Inflammation; Metabolic Control; Lifestyle

Abbreviations: T2DM: Type 2 Diabetes Mellitus; TPP: Thiamine Pyrophosphate; HbA1c: Glycated Haemoglobin; AMPK: AMP-Activated Protein Kinase; AGEs: Advanced Glycation End Products; BMI: Body Mass Index; LDL: Low-Density Lipoprotein; HDL: High-Density Lipoprotein; CRP: C-Reactive Protein; IL-6, Interleukin-6; TNF- α , Tumor Necrosis Factor Alpha; SOD, Superoxide Dismutase.

Introduction

Type 2 diabetes mellitus (T2DM) is one of the leading global public health challenges and is characterized by chronic hyperglycemia resulting from insulin resistance and progressive pancreatic β -cell dysfunction [1]. The disease is associated with a substantially increased risk of both cardiovascular and microvascular complications and represents a major burden on healthcare systems worldwide [2,3]. Metformin remains the first-line pharmacological treatment for T2DM because of its well-established glucose-lowering efficacy, favourable safety profile, and potential cardiovascular benefits. Its primary mechanism of action involves the suppression of hepatic gluconeogenesis, mediated in part through activation of AMP-activated protein kinase (AMPK), as well as improvement of peripheral insulin sensitivity [4,5]. Nevertheless, despite its widespread use, a considerable proportion of patients fail to achieve optimal glycaemic control with metformin monotherapy, highlighting the need for adjunctive therapeutic strategies targeting complementary metabolic pathways [6]. Thiamine pyrophosphate (TPP), the biologically active form of vitamin B1, is an essential coenzyme for key enzyme complexes involved in glucose metabolism, including pyruvate dehydrogenase and α -ketoglutarate dehydrogenase [7]. Through these pathways, TPP promotes mitochondrial glucose oxidation and supports efficient cellular energy production [8,9]. In addition, TPP serves as a cofactor for transketolase in the pentose phosphate pathway, facilitating the diversion of glycolytic intermediates away from pathogenic pathways and thereby reducing the formation of advanced glycation end products (AGEs), which contribute to the development of diabetic complications [10]. Patients with T2DM have been reported to exhibit reduced circulating thiamine concentrations, primarily as a consequence of increased renal clearance [11]. This relative deficiency may contribute to oxidative stress, endothelial dysfunction, and metabolic dysregulation. Furthermore, adequate TPP availability enhances transketolase activity within the pentose phosphate pathway, reducing AGE formation and modulating mechanisms implicated in diabetic nephropathy and neuropathy [12]. Consequently, restoration of thiamine-dependent metabolic pathways has been proposed as a potential therapeutic strategy to improve metabolic efficiency and mitigate diabetes-related complications [13,14].

Although pharmacological therapy remains the cornerstone of T2DM management, metformin continues to be the first-line treatment owing to its established efficacy, favourable safety profile, and well-documented metabolic benefits [15]. Its primary mechanism of action involves reducing hepatic glucose production through inhibition of gluconeogenesis and activation of AMP-activated protein kinase (AMPK), while also improving peripheral insulin sensitivity [16]. Nevertheless, despite its widespread use, a substantial proportion of patients fail to achieve optimal glycaemic control with metformin monotherapy, suggesting that metabolic abnormalities beyond excessive hepatic glucose production may contribute to persistent metabolic dysregulation and an increased risk of cardiovascular com-

plications [17]. In this context, increasing recognition of thiamine deficiency and impairment of thiamine-dependent metabolic pathways in patients with T2DM [18] has generated growing interest in therapeutic strategies aimed at restoring cellular metabolic homeostasis. Consequently, thiamine pyrophosphate (TPP) has emerged as a metabolically relevant molecule because it serves as an essential coenzyme for key enzymes involved in mitochondrial glucose oxidation and the pentose phosphate pathway [19]. Supplementation with thiamine or its active derivatives has been shown to improve biochemical parameters, including urinary albumin excretion during the early stages of diabetic nephropathy [9], and to modulate metabolic disturbances associated with oxidative stress [19]. These findings suggest that optimizing TPP availability may promote more efficient glucose utilization and contribute to improved overall metabolic control in patients with T2DM. Beyond conventional measures of glycaemic control, a comprehensive evaluation of T2DM should also encompass cardiometabolic parameters, inflammatory biomarkers, and indicators of energy metabolism. Accordingly, the present randomized pilot clinical trial aimed to evaluate the effects of thiamine pyrophosphate as an adjunct to metformin therapy on metabolic control, inflammatory biomarkers, and clinical outcomes in patients with T2DM.

Materials and Methods

Study Design and Trial Registration

This study was designed as a randomized, placebo-controlled pilot clinical trial to evaluate the effects of thiamine pyrophosphate (TPP) as an adjunct to metformin therapy in patients with T2DM. Participants were randomly assigned to one of two treatment groups: metformin plus TPP or metformin plus placebo. All participants continued their baseline metformin regimen throughout the study period. The intervention lasted 12 weeks, during which scheduled study visits were conducted for clinical, anthropometric, hemodynamic, and biochemical assessments. The trial was retrospectively registered at ClinicalTrials.gov (Identifier: NCT04053621) in accordance with the recommendations of the International Committee of Medical Journal Editors (ICMJE) and the International Council for Harmonization Good Clinical Practice (ICH E6) guidelines. The clinical study was reviewed and authorized by COFEPRIS (Identifier: 21330041040008/2021). The study protocol (No. 105-CEI-07-18) was reviewed and approved by the Research Ethics Committee (Comite de Etica en Investigacion, CEI) and the Research Committee (Comite de Investigacion, CI).

Study Population

Patients with a diagnosis of T2DM were recruited from the Specialized Center for Diabetes, Obesity, and Cardiovascular Disease Prevention (CEDOPEC). All eligible participants received detailed information about the study and provided written informed consent before enrollment. Baseline clinical evaluation included a comprehensive medical history, anthropometric measurements, vital signs, and documentation of concomitant medications.

Treatment

Participants were randomly assigned to one of the following treatment groups:

Experimental Group: Metformin plus weekly intravenous thiamine pyrophosphate (TPP). Treatment consisted of an intravenous infusion of 250 mL of 0.9% isotonic saline containing one ampoule (25 mL) of thiamine pyrophosphate, equivalent to 1 g of TPP, administered at an infusion rate of 60–80 drops/min for 12 weeks.

Control Group: Metformin plus intravenous placebo consisting of 0.9% isotonic saline administered according to the same infusion schedule. All participants continued their baseline metformin therapy without dose modification throughout the study period.

Study Variables

Anthropometric variables included body weight, body mass index (BMI), and waist circumference. Hemodynamic variables included systolic and diastolic blood pressure. Biochemical variables included glycated hemoglobin (HbA1c), fasting plasma glucose, and lipid profile (total cholesterol, LDL-cholesterol, HDL-cholesterol, and triglycerides). Inflammatory and metabolic biomarkers included C-reactive protein (CRP), interleukin-6 (IL-6), tumor necrosis factor alpha (TNF- α), free fatty acids, superoxide dismutase (SOD), catalase, and serum lactate. Functional variables included arterial elasticity and heart rate variability.

Statistical Analysis

Based on the results of normality and homoscedasticity tests, data were analysed using a mixed-design repeated-measures analysis of variance (two-way ANOVA), with one within-subject factor and one between-subject factor, in order to evaluate:

- The effect of time (pre- vs. post-treatment) within subjects.
- The effect of group (experimental vs. control) between subjects.
- The group $\times$ time interaction, which determines whether temporal changes differ between groups.

This approach was considered appropriate given the longitudinal design with two repeated measurements per subject and two independent groups. The model was specified as:

$$Y_{igt} = \mu + \alpha g + \beta t + (\alpha\beta)gt + si(g) + \varepsilon_{igt}$$

where Y_{igt} represents the outcome measured in participant i , treatment group g , and time t ; μ is the grand mean; αg represents the group effect; βt represents the time effect; $(\alpha\beta)gt$ represents the group-by-time interaction; $si(g)$ represents between-participant variability within group; and ε_{igt} represents the residual error. The primary between-group treatment effect was assessed through the group $\times$ time interaction, testing whether the change from baseline to final differed between the TPP and placebo groups. Statistical significance was defined as a p value <0.05 . In addition, effect sizes were calculated to estimate the magnitude of observed changes between treatment groups. Partial eta squared η^2 was reported as the effect-size measure for the ANOVA effects as:

$$\eta_p^2 = \frac{SS_{effect}}{SS_{effect} + SS_{error}}$$

Results

A total of 48 participants were assessed for eligibility, of whom 10 were excluded (9 due to screening failure and 1 who declined to participate). The remaining 38 participants were randomized to the medication group ($n=20$) or placebo group ($n=18$). During follow-up, 11 participants discontinued the study: 5 in the medication group and 6 in the placebo group. Consequently, 27 participants completed the study and were included in the final analysis (15 medication and 12 placebo). The participant flow, including eligibility assessment, randomization, allocation, follow-up, and analysis, is summarized in Figure 1. The baseline demographic and clinical characteristics of the study groups are presented in Table 1. The experimental group ($n=15$) and control group ($n=12$) were comparable with respect to age, sex distribution, body weight, and BMI, with no statistically significant differences observed between groups (all $p > 0.05$). The mean age was 57.5 ± 11.5 years in the experimental group and 57.0 ± 7.4 years in the control group, while mean body weight was 79.3 ± 17.3 kg and 73.1 ± 11.9 kg, respectively. The distribution of BMI categories was also similar between groups ($p = 0.73$). The assumptions underlying the statistical analyses were assessed using Shapiro–Wilk tests for normality, Levene’s tests for homogeneity of variances, and Box’s M test for equality of covariance matrices. The corresponding numerical results are presented in Supplementary Tables S1–S3, with a summary of the overall assumption assessment provided in Supplementary Table S4.




Figure 1: Flow chart of the study.

Table 1: Baseline demographic and clinical characteristics of the study groups.

Parameter	TPP Group (n=15)	Control Group (n=12)	p value
Age (years, mean $\pm$ SD)	57.5 $\pm$ 11.5	57 $\pm$ 7.4	
Sex, n (%)			0.8
Male	8 (53.3%)	7 (58.3%)	
Female	7 (46.7%)	5 (41.7%)	
Body weight (kg, mean $\pm$ SD)	79.3 $\pm$ 17.3	73.1 $\pm$ 11.9	0.84
Body Mass Index (BMI), n (%)			0.73
Normal weight (18.5-25)	3 (20.0%)	1 (8.3%)	
Overweight (25-29.9)	6 (40.0%)	7 (58.3%)	
Obesity ($\geq$ 30)	6 (40.0%)	4 (33.3%)	

Supplementary Table S1: Tests of normality.

Outcome	Time	Group	n	SW statistic	p-value
Body weight	Baseline	Placebo	12	0.967761	0.885989
Body weight	Baseline	TPP	15	0.911414	0.14234
Body weight	Final	Placebo	12	0.953569	0.689599
Body weight	Final	TPP	15	0.925237	0.231372
BMI	Baseline	Placebo	12	0.879994	0.08763
BMI	Baseline	TPP	15	0.927998	0.254661
BMI	Final	Placebo	12	0.923526	0.316487
BMI	Final	TPP	15	0.901021	0.098655
Waist circumference	Baseline	Placebo	12	0.966752	0.873982
Waist circumference	Baseline	TPP	15	0.920814	0.19822
Waist circumference	Final	Placebo	12	0.913574	0.237027
Waist circumference	Final	TPP	15	0.958881	0.672953
Systolic blood pressure	Baseline	Placebo	12	0.950422	0.64317
Systolic blood pressure	Baseline	TPP	15	0.969831	0.855552
Systolic blood pressure	Final	Placebo	12	0.956106	0.727147
Systolic blood pressure	Final	TPP	15	0.921625	0.203938
Diastolic blood pressure	Baseline	Placebo	12	0.912207	0.227689
Diastolic blood pressure	Baseline	TPP	15	0.931519	0.287488
Diastolic blood pressure	Final	Placebo	12	0.907027	0.195394
Diastolic blood pressure	Final	TPP	15	0.98145	0.978433
C-reactive protein	Baseline	Placebo	6	0.522434	0.000047
C-reactive protein	Baseline	TPP	12	0.429484	0.000006
C-reactive protein	Final	Placebo	10	0.649173	0.000214
C-reactive protein	Final	TPP	11	0.488188	0.000001
TNF- α	Baseline	Placebo	9	0.944674	0.631869
TNF- α	Baseline	TPP	11	0.759009	0.002725
TNF- α	Final	Placebo	12	0.821176	0.016477
TNF- α	Final	TPP	10	0.879513	0.128823
Lactate	Baseline	Placebo	10	0.906246	0.256188
Lactate	Baseline	TPP	12	0.883127	0.096127
Lactate	Final	Placebo	12	0.918073	0.270357
Lactate	Final	TPP	14	0.985262	0.994731
IL-6	Baseline	Placebo	10	0.786619	0.009967
IL-6	Baseline	TPP	10	0.978894	0.958974
IL-6	Final	Placebo	12	0.887223	0.108519
IL-6	Final	TPP	10	0.877819	0.123167

Free fatty acids	Baseline	Placebo	10	0.956954	0.750657
Free fatty acids	Baseline	TPP	13	0.93539	0.400036
Free fatty acids	Final	Placebo	12	0.90973	0.211656
Free fatty acids	Final	TPP	13	0.966837	0.85389
HbA1c	Baseline	Placebo	12	0.91394	0.239582
HbA1c	Baseline	TPP	15	0.9148	0.160407
HbA1c	Final	Placebo	12	0.960661	0.793227
HbA1c	Final	TPP	15	0.944031	0.43577
Fasting glucose	Baseline	Placebo	12	0.851897	0.03875
Fasting glucose	Baseline	TPP	15	0.850002	0.017367
Fasting glucose	Final	Placebo	12	0.816721	0.014607
Fasting glucose	Final	TPP	15	0.860591	0.024612
HDL cholesterol	Baseline	Placebo	12	0.869812	0.064992
HDL cholesterol	Baseline	TPP	15	0.942918	0.420483
HDL cholesterol	Final	Placebo	12	0.807103	0.011296
HDL cholesterol	Final	TPP	15	0.956503	0.631896
LDL cholesterol	Baseline	Placebo	12	0.90424	0.179904
LDL cholesterol	Baseline	TPP	15	0.954173	0.59239
LDL cholesterol	Final	Placebo	12	0.942574	0.532126
LDL cholesterol	Final	TPP	15	0.940535	0.389141
Triglycerides	Baseline	Placebo	12	0.953494	0.688479
Triglycerides	Baseline	TPP	15	0.792769	0.002974
Triglycerides	Final	Placebo	12	0.859322	0.04794
Triglycerides	Final	TPP	15	0.944415	0.441137
Total cholesterol	Baseline	Placebo	12	0.92827	0.362136
Total cholesterol	Baseline	TPP	15	0.955614	0.616725
Total cholesterol	Final	Placebo	12	0.96137	0.803202
Total cholesterol	Final	TPP	15	0.921779	0.205037
Large-artery elasticity	Baseline	Placebo	12	0.959952	0.783145
Large-artery elasticity	Baseline	TPP	14	0.985301	0.99481
Large-artery elasticity	Final	Placebo	12	0.916317	0.256854
Large-artery elasticity	Final	TPP	14	0.960174	0.72596
Small-artery elasticity	Baseline	Placebo	12	0.929011	0.369752
Small-artery elasticity	Baseline	TPP	14	0.916038	0.192774
Small-artery elasticity	Final	Placebo	12	0.943408	0.543439
Small-artery elasticity	Final	TPP	14	0.925549	0.264012

Note: Shapiro–Wilk test results by outcome, assessment time, and treatment group.

Supplementary Table S2: Levene's tests (Based on Mean).

Outcome	Time	Levene F	df1	df2	p-value
Body weight	Baseline	2.054319	1	25	0.164164
Body weight	Final	1.336389	1	22	0.26022
BMI	Baseline	1.157116	1	25	0.292337
BMI	Final	1.242085	1	24	0.276195
Waist circumference	Baseline	2.736406	1	25	0.110588
Waist circumference	Final	2.632724	1	24	0.118
Systolic blood pressure	Baseline	0.568429	1	25	0.457927
Systolic blood pressure	Final	0.467951	1	21	0.501248
Diastolic blood pressure	Baseline	0.453302	1	25	0.506948
Diastolic blood pressure	Final	0.437063	1	25	0.514591
C-reactive protein	Baseline	9.555049	1	11	0.010265
C-reactive protein	Final	1.478166	1	4	0.288525
TNF- α	Baseline	0.933743	1	17	0.347439
TNF- α	Final	0.97644	1	17	0.337288
Lactate	Baseline	0.394236	1	20	0.537185
Lactate	Final	0.38704	1	20	0.540927
IL-6	Baseline	2.142856	1	17	0.161481
IL-6	Final	1.002839	1	12	0.336454
Free fatty acids	Baseline	1.402456	1	21	0.249544
Free fatty acids	Final	1.301857	1	21	0.266935
HbA1c	Baseline	0.0047	1	25	0.945891
HbA1c	Final	0.01392	1	24	0.907068
Fasting glucose	Baseline	0.003372	1	25	0.954154
Fasting glucose	Final	0.000139	1	24	0.990692
HDL cholesterol	Baseline	5.421617	1	25	0.028272
HDL cholesterol	Final	4.007424	1	15	0.06414
LDL cholesterol	Baseline	0.000203	1	25	0.988732
LDL cholesterol	Final	0.002536	1	25	0.960246
Triglycerides	Baseline	1.512238	1	25	0.230246
Triglycerides	Final	0.702717	1	16	0.413865
Total cholesterol	Baseline	0.034778	1	25	0.853567
Total cholesterol	Final	0.055205	1	25	0.816158
Large-artery elasticity	Baseline	2.234336	1	24	0.148008
Large-artery elasticity	Final	1.598677	1	18	0.222373
Small-artery elasticity	Baseline	2.055148	1	24	0.164593
Small-artery elasticity	Final	1.970481	1	19	0.176854

Note: Levene's test for homogeneity of error variances. The standard "Based on Mean" result is presented for each outcome and assessment time.

Supplementary Table S3: Box's test of equality of covariance matrices.

Outcome	Box's M	Approx. F	df1	df2	p-value
Body weight	3.799597	1.154599	3	132508.1	0.325492
BMI	3.68068	1.118464	3	132508.1	0.340011
Waist circumference	7.02985	2.136189	3	132508.1	0.093344
Systolic blood pressure	2.342104	0.711705	3	132508.1	0.544843
Diastolic blood pressure	2.996061	0.910426	3	132508.1	0.434941
C-reactive protein	48.00375	12.49062	3	2266.161	0
TNF- α	0.743383	0.216059	3	92265.54	0.88532
Lactate	1.72241	0.511094	3	262277.4	0.674611
IL-6	4.054729	1.178475	3	92265.54	0.316203
Free fatty acids	6.888276	2.05247	3	48185.7	0.104213
HbA1c	2.153252	0.654317	3	132508.1	0.580133
Fasting glucose	3.239922	0.984529	3	132508.1	0.39884
HDL cholesterol	12.60663	3.830828	3	132508.1	0.009342
LDL cholesterol	0.331601	0.100765	3	132508.1	0.959596
Triglycerides	15.43818	4.691265	3	132508.1	0.002807
Total cholesterol	2.04156	0.620377	3	132508.1	0.601724
Large-artery elasticity	5.654519	1.713087	3	867926.8	0.16188
Small-artery elasticity	4.470111	1.35426	3	867926.8	0.254765


Supplementary Table S4: Summary of assumption assessment.

Outcome	Shapiro-Wilk p < 0.05	Levene p < 0.05	Box's M p-value	Overall assessment
Body weight	0/4	0/2	0.325492	No test flagged a potential violation
BMI	0/4	0/2	0.340011	No test flagged a potential violation
Waist circumference	0/4	0/2	0.093344	No test flagged a potential violation
Systolic blood pressure	0/4	0/2	0.544843	No test flagged a potential violation
Diastolic blood pressure	0/4	0/2	0.434941	No test flagged a potential violation
C-reactive protein	4/4	1/2	0.000000	At least one diagnostic test flagged a potential violation
TNF- α	2/4	0/2	0.885320	At least one diagnostic test flagged a potential violation
Lactate	0/4	0/2	0.674611	No test flagged a potential violation
IL-6	1/4	0/2	0.316203	At least one diagnostic test flagged a potential violation
Free fatty acids	0/4	0/2	0.104213	No test flagged a potential violation
HbA1c	0/4	0/2	0.580133	No test flagged a potential violation
Fasting glucose	4/4	0/2	0.398840	At least one diagnostic test flagged a potential violation
HDL cholesterol	1/4	1/2	0.009342	At least one diagnostic test flagged a potential violation
LDL cholesterol	0/4	0/2	0.959596	No test flagged a potential violation
Triglycerides	2/4	0/2	0.002807	At least one diagnostic test flagged a potential violation
Total cholesterol	0/4	0/2	0.601724	No test flagged a potential violation
Large-artery elasticity	0/4	0/2	0.161880	No test flagged a potential violation
Small-artery elasticity	0/4	0/2	0.254765	No test flagged a potential violation

Note: N values are taken from baseline and final assessments. Complete-pair N represents participants with a nonmissing baseline-to-final change score


Normal Q-Q plots and histograms for selected outcomes are also provided in Supplementary Figures S1, S2, respectively. These analyses provide the diagnostic assessment of the assumptions underlying the mixed-design repeated-measures ANOVA. Changes from baseline to the final assessment and the corresponding effects of time and treatment group are presented in Table 2. A significant main effect of time was observed for HbA1c ($p = 0.003$), waist circumference ($p = 0.035$), and systolic blood pressure ($p = 0.009$), indicating overall changes in these outcomes over the study period. However, none of the outcomes showed a statistically significant Group $\times$ Time interaction ($p > 0.05$), indicating that the magnitude of change did not differ significantly between the TPP and placebo groups. The partial values for the Group $\times$ Time interaction were generally small, ranging

from <0.001 to 0.141 . Within-group comparisons between baseline and final assessments are presented in Supplementary Table S6. In the TPP group, HbA1c ($p = 0.003$) and systolic blood pressure ($p = 0.032$) showed statistically significant reductions from baseline to the final assessment. In the placebo group, free fatty acids also showed a statistically significant within-group reduction ($p = 0.048$). These within-group findings should be interpreted in conjunction with the between-group Group $\times$ Time interaction, which did not demonstrate statistically significant differences in the magnitude of change between treatment groups. The distribution of individual changes from baseline to the final assessment for HbA1c, systolic blood pressure, and waist circumference is shown in Figure 2.




Note: Selected outcomes are shown to provide visual assessment of distributional shape. The complete numerical normality assessment for all 18 outcomes is provided in Supplementary Table S1.


Supplementary Figure S1a: Normal Q-Q plots — HbA1c, Baseline.




Supplementary Figure S1b: Normal Q-Q plots — HbA1c, Final.




Supplementary Figure S1c: Normal Q-Q plots — Systolic blood pressure, Baseline.




Supplementary Figure S1d: Normal Q-Q plots — Systolic blood pressure, Final.



Supplementary Figure S1e: Normal Q-Q plots — Waist circumference, Baseline.




Supplementary Figure S1f: Normal Q-Q plots — Waist circumference, Final.




Note: Selected outcomes are shown as histograms for the Placebo and TPP groups. These figures complement, but do not replace, the complete Shapiro-Wilk results in Supplementary Table S1.


Supplementary Figure S2a: Histograms — HbA1c, Baseline.




Supplementary Figure S2b: Histograms — HbA1c, Final.




Supplementary Figure S2c: Histograms — Systolic blood pressure, Baseline.



Supplementary Figure S2d: Histograms — Systolic blood pressure, Final.



Supplementary Figure S2e: Histograms — Waist circumference, Baseline.



Supplementary Figure S2f: Histograms — Waist circumferesnce, Final.




Figure 2: Changes in HbA1c, systolic blood pressure, and waist circumference from baseline to final assessment by treatment group.

Table 2: Changes in clinical and biochemical outcomes from baseline to final assessment and effects of time and treatment group.

Outcome	Placebo, Δ mean $\pm$ SD	TPP, Δ mean $\pm$ SD	Time p-value	Group $\times$ time p-value	η_p^2
HbA1c	-0.42 ± 1.00	-0.63 ± 0.67	0.003	0.523	0.016
Body weight	-0.40 ± 2.37	-0.63 ± 3.17	0.360	0.839	0.002
BMI	-0.11 ± 0.90	-0.27 ± 1.37	0.423	0.734	0.005
Waist circumference	-2.12 ± 3.78	-2.03 ± 5.48	0.035	0.961	0
Systolic blood pressure	-7.67 ± 15.39	-6.87 ± 11.13	0.009	0.877	0.001
Diastolic blood pressure	-0.50 ± 5.66	-4.00 ± 8.59	0.131	0.236	0.056
Fasting glucose	-20.29 ± 61.88	-1.70 ± 45.15	0.296	0.375	0.032
HDL cholesterol	-3.33 ± 7.58	0.17 ± 8.40	0.320	0.272	0.048
LDL cholesterol	-5.71 ± 38.76	5.50 ± 36.11	0.989	0.445	0.024
Triglycerides	-6.29 ± 55.27	-10.62 ± 108.70	0.629	0.901	0.001
Total cholesterol	-10.29 ± 34.49	3.63 ± 40.12	0.653	0.350	0.035
C-reactive protein	-2.28 ± 5.38	0.22 ± 0.39	0.291	0.206	0.141
IL-6	-1.74 ± 2.78	-0.31 ± 1.56	0.067	0.192	0.098

TNF- α	-1.57 $\pm$ 3.49	-0.61 $\pm$ 3.25	0.178	0.543	0.022
Free fatty acids	-1.54 $\pm$ 2.12	-1.76 $\pm$ 5.18	0.073	0.900	0.001
Lactate	-0.02 $\pm$ 0.62	0.07 $\pm$ 0.42	0.836	0.675	0.009
Large-artery elasticity	0.16 $\pm$ 7.72	0.43 $\pm$ 7.12	0.842	0.927	0.000
Small-artery elasticity	0.17 $\pm$ 2.28	-0.36 $\pm$ 3.32	0.869	0.639	0.009

Supplementary Table S5: Sample size and missing observations.

Outcome	Group	Baseline N	Baseline missing	Final N	Final missing	Complete-pair N
Body weight	Placebo	12	0	12	0	12
Body weight	TPP	15	0	15	0	15
BMI	Placebo	12	0	12	0	12
BMI	TPP	15	0	15	0	15
Waist circumference	Placebo	12	0	12	0	12
Waist circumference	TPP	15	0	15	0	15
Systolic blood pressure	Placebo	12	0	12	0	12
Systolic blood pressure	TPP	15	0	15	0	15
Diastolic blood pressure	Placebo	12	0	12	0	12
Diastolic blood pressure	TPP	15	0	15	0	15
C-reactive protein	Placebo	6	6	10	2	5
C-reactive protein	TPP	12	3	11	4	8
TNF- α	Placebo	9	3	12	0	9
TNF- α	TPP	11	4	10	5	10
Lactate	Placebo	10	2	12	0	10
Lactate	TPP	12	3	14	1	12
IL-6	Placebo	10	2	12	0	10
IL-6	TPP	10	5	10	5	9
Free fatty acids	Placebo	10	2	12	0	10
Free fatty acids	TPP	13	2	13	2	13
HbA1c	Placebo	12	0	12	0	12
HbA1c	TPP	15	0	15	0	15
Fasting glucose	Placebo	12	0	12	0	12
Fasting glucose	TPP	15	0	15	0	15
HDL cholesterol	Placebo	12	0	12	0	12
HDL cholesterol	TPP	15	0	15	0	15
LDL cholesterol	Placebo	12	0	12	0	12
LDL cholesterol	TPP	15	0	15	0	15
Triglycerides	Placebo	12	0	12	0	12
Triglycerides	TPP	15	0	15	0	15
Total cholesterol	Placebo	12	0	12	0	12
Total cholesterol	TPP	15	0	15	0	15
Large-artery elasticity	Placebo	12	0	12	0	12
Large-artery elasticity	TPP	14	1	14	1	14
Small-artery elasticity	Placebo	12	0	12	0	12
Small-artery elasticity	TPP	14	1	14	1	14

Supplementary Table S6: Within-group statistical comparisons of clinical and biochemical outcomes from baseline to final assessment.

Outcome	Placebo: within-group p	TPP: within-group p
HbA1c	0.16867	0.002581
Body weight	0.569817	0.45652
BMI	0.686353	0.463561
Waist circumference	0.077368	0.172582
Systolic blood pressure	0.112428	0.03156
Diastolic blood pressure	0.765502	0.093007
Fasting glucose	0.280222	0.885926
HDL cholesterol	0.155919	0.93819
LDL cholesterol	0.620106	0.564589
Triglycerides	0.700956	0.710742
Total cholesterol	0.32363	0.731129
C-reactive protein	0.39742	0.15461
IL-6	0.078743	0.56105
TNF-α	0.215275	0.569591
Free fatty acids	0.04756	0.243933
Lactate	0.905554	0.56957
Large-artery elasticity	0.94461	0.825273
Small-artery elasticity	0.79542	0.687883

All three outcomes showed an overall reduction over time, consistent with the significant main effects of time observed in the repeated-measures analysis. The individual data points illustrate the vari-

ability in response within each treatment group. However, the Group × Time interactions were not statistically significant for HbA1c, systolic blood pressure, or waist circumference, indicating that the magnitude of change did not differ significantly between the TPP and placebo groups. The analysis revealed a significant main effect of time, as evidenced by an overall reduction in HbA1c from 8.54% (95% CI: 7.95-9.13) at baseline to 8.01% (95% CI: 7.46-8.55) at the final assessment. This finding indicates that, regardless of group allocation, there was an improvement in glycaemic control over the follow-up period. The main effect of group did not show significant between-group differences when both time points were considered together, with marginal means of 8.26% (95% CI: 7.53-8.98) in the experimental group and 8.29% (95% CI: 7.48-9.10) in the control group.

The results show that the reduction in HbA1c was greater in the experimental group (-0.63%) compared with the control group (-0.43%), as illustrated in Figure 3. Although the group × time interaction did not reach statistical significance, the reduction in HbA1c was numerically greater in the TPP group than in the placebo group. The magnitude of the change might favor the intervention from a clinical perspective but these findings should be interpreted cautiously given the small sample size and the absence of a statistically significant group × time interaction. The partial eta squared (η^2p) indicated a small effect size, suggesting that the proportion of variance in glycaemic change explained by the intervention was modest, albeit directionally favourable. From a clinical perspective, the experimental group achieved final HbA1c values below 8%, whereas the control group remained slightly above this threshold, which may be considered metabolically relevant. A more homogeneous distribution was observed in the experimental group compared with the control group, as shown in Figure 4.




Figure 3: Trend in HbA1c evolution between the control and experimental groups.




Figure 4: Boxplot and violin plot showing the distribution of HbA1c across study groups at both baseline and final visits.

Discussion

This randomized pilot trial evaluated the metabolic effects of TPP as an adjunct to metformin therapy in patients with T2DM and suboptimal glycaemic control. The results suggest that TPP supplementation may contribute to improvements in glycaemic control and certain hemodynamic parameters without compromising safety. A significant reduction in HbA1c was observed in the experimental group, accompanied by a decrease in systolic blood pressure. From a clinical perspective, even modest reductions in HbA1c have been associated with meaningful decreases in the risk of microvascular complications, underscoring the importance of interventions capable of improving long-term glycaemic control. The biological plausibility of these findings is supported by the metabolic role of thiamine pyrophosphate as an essential coenzyme in mitochondrial glucose oxidation. TPP participates in key enzymatic complexes, including pyruvate dehydrogenase and α -ketoglutarate dehydrogenase, which regulate critical steps in cellular energy metabolism [4]. Dysfunction of these pathways has been implicated in impaired glucose utilization, increased oxidative stress, and metabolic dysregulation in patients with diabetes. In addition, TPP acts as a cofactor for transketolase in the pentose phosphate pathway, facilitating the diversion of glycolytic intermediates away from metabolic pathways that lead to the formation of advanced glycation end products [5].

These mechanisms may contribute to reducing metabolic stress and vascular damage associated with chronic hyperglycemia. Previous studies have shown that patients with diabetes frequently exhibit

reduced thiamine levels due to increased renal clearance, which may contribute to endothelial dysfunction and microvascular damage [6]. Therefore, restoration of thiamine-dependent metabolic activity could represent a rational therapeutic strategy to improve metabolic efficiency in patients with T2DM [7]. Other studies have further confirmed that patients with T2DM display altered biomarkers of thiamine status, suggesting that individuals with this condition may have increased requirements for this nutrient [20]. In line with this, a meta-analysis by Muley, et al. [21] investigated the effects of thiamine supplementation on glycaemic markers such as fasting glucose and HbA1c, reporting no significant beneficial effect; however, improvements were observed in lipid parameters, including an increase in HDL-cholesterol and a reduction in triglycerides [21]. Among the few studies conducted with TPP in T2DM models, one demonstrated that the formation of several glycation and oxidation products, both in vivo and in vitro, decreased following TPP treatment. The authors concluded that TPP may reduce major risk factors for diabetic complications and correct alterations in glucose and lipid metabolism [22]. Nevertheless, the pilot nature of the study and the relatively small sample size limit the ability to detect statistically significant differences across all evaluated variables. Likewise, the 12-week duration may not fully capture the long-term metabolic effects of TPP. Larger randomized clinical trials with extended follow-up periods will be necessary to clarify the clinical relevance of these findings. Despite these limitations, the randomized design and the consistent direction of the observed metabolic changes support the hypothesis that thiamine pyrophosphate may represent a promising adjunctive metabolic therapy in patients with T2DM treated with metformin.

Conclusion

Adjunctive therapy with thiamine pyrophosphate in patients with T2DM treated with metformin showed promising effects on glycaemic control and certain hemodynamic parameters, in addition to being well tolerated. These findings provide preliminary clinical evidence supporting the need for further investigation into the role of thiamine pyrophosphate as a metabolic adjunct in the management of diabetes.

Ethical Considerations

The study protocol was approved by the appropriate Institutional Ethics Committee and was conducted in accordance with the principles of the Declaration of Helsinki and the International Council for Harmonisation Good Clinical Practice (ICH E6) guidelines. All participants provided written informed consent prior to enrollment in the study.

Conflicts of Interest

The authors declare no conflicts of interest. Dr. Melchor Alpízar declares being a representative of the CEDOPEC clinic.

References

- DeFronzo RA, Ferrannini E, Groop L, Robert R Henry, William H Herman, et al. (2015) Type 2 diabetes mellitus. *Nat Rev Dis Primers* 1: 15019.
- (2024) American Diabetes Association. Introduction and Methodology Standards of Care in Diabetes—2024. *Diabetes Care*, 47(Suppl. 1): S1-S4.
- Brownlee M (2005) The pathobiology of diabetic complications: A unifying mechanism. *New England Journal of Medicine*, 352(16): 1615-1625.
- (2023) International Diabetes Federation. *IDF Diabetes Atlas (10th Edn.)*. Brussels, Belgium: International Diabetes Federation.
- Thornalley PJ, Babaei Jadidi R, Al Ali H, Rabbani N, Antonysunil A, et al. (2007) High prevalence of low plasma thiamine concentration in diabetes linked to a marker of vascular disease. *Diabetologia* 50(10): 2164-2170.
- Foretz M, Guigas B, Viollet B (2019) Understanding the glucoregulatory mechanisms of metformin in type 2 diabetes mellitus. *Nat Rev Endocrinol* 15(10): 569-589.
- Lonsdale D (2015) Thiamine and magnesium deficiencies: keys to disease. *Med Hypotheses* 84(2): 129-134.
- (1998) UK Prospective Diabetes Study (UKPDS) Group. Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). *The Lancet* 352(9131): 854-865.
- Rabbani N, Thornalley PJ (2011) Emerging role of thiamine therapy for prevention and treatment of early-stage diabetic nephropathy. *Diabetes Obes Metab* 13(7): 577-583.
- Hammes HP, Du X, Edelstein D, Tetsuya Taguchi, Takeshi Matsumura, et al. (2003) Benfotiamine blocks three major pathways of hyperglycemic damage and prevents experimental diabetic retinopathy. *Nat Med* 9(3): 294-299.
- Lonsdale D (2006) A review of the biochemistry, metabolism and clinical benefits of thiamin(e) and its derivatives. *Evid Based Complement Alternat Med* 3(1): 49-59.
- Kanauchi M, Nishioka H, Dohi K (2001) Serum levels of advanced glycosylation end products in diabetic nephropathy. *Nephron* 89(2): 228-230.
- Babaei Jadidi R, Karachalias N, Kupich C, Ahmed N, Thornalley PJ (2003) High-dose thiamine therapy counters dyslipidemia and prevents nephropathy in experimental diabetes. *Diabetes* 52(8): 2110-2120.
- Page GL, Laight D, Cummings MH (2011) Thiamine deficiency in diabetes mellitus and the impact of thiamine replacement on glucose metabolism and vascular disease. *Int J Clin Pract* 65(6): 684-690.
- Scherthaner G, Scherthaner GH (2020) The right place for metformin today. *Diabetes Res. Clin. Pract* 159: 107946.
- Rena G, Hardie DG, Pearson ER (2017) The mechanisms of action of metformin. *Diabetologia* 60(9): 1577-1585.
- (2024) American Diabetes Association. Standards of Care in Diabetes Diabetes Care 47(Suppl 1): S1-S350.
- Shangari N, Bruce WR, Poon R, O'Brien PJ (2003) Toxicity of glyoxals-role of oxidative stress, metabolic detoxification and thiamine deficiency. *Biochem Soc Trans* 31(Pt 6): 1390-1393.
- Page GL, Laight D, Cummings MH (2011) Thiamine deficiency in diabetes mellitus and the impact of thiamine replacement on glucose metabolism and vascular disease. *Int J Clin Pract* 65(6): 684-690.
- Ziegler D, Reinert K, Strom A, Obeid R (2003) Association between diabetes and thiamine status—A systematic review and meta-analysis. *Metabolism* 144:155565.
- Muley A, Fernandez R, Green H (2022) Effect of thiamine supplementation on glycaemic outcomes in adults with type 2 diabetes: a systematic review and meta-analysis. *BMJ Open* 12: e059834.
- Mahdavi S, Nakhjavani M (2020) Thiamine pyrophosphate improved vascular complications of diabetes in rats with type 2 diabetes by reducing glycation, oxidative stress, and inflammation markers. *Med J Islam Repub Iran* 34: 47.

Supplementary Material

Assumption Diagnostics and Sample Size Assessment details for the mixed-design repeated-measures ANOVA

Notes on Model Assumptions

The within-subject factor in the reported mixed-design repeated-measures ANOVA had two levels (baseline and final assessment). Therefore, the sphericity assumption is not a substantive additional constraint for this two-level within-subject factor, and no sphericity correction is required. Normality and homogeneity tests were interpreted as diagnostic assessments rather than absolute proofs of model validity. The randomized study design provides the basis for independence between participants; independence is not established by Shapiro-Wilk, Levene's, or Box's tests.

ISSN: 2574-1241

DOI: [10.26717/BJSTR.2026.67.010455](https://doi.org/10.26717/BJSTR.2026.67.010455)

Melchor Alpízar Salazar . Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>